

Abstract

Relationship between the Use of Mobile Phones and
Hearing Thresholds in Some White-collar Workers

Mi Young Lee, Choong Won Lee

Department of Preventive Medicine and Institute for Medical Science
College of Medicine, Keimyung University

Objectives: The purpose of this study was to investigate the relationship between the use of mobile phones and the levels of hearing thresholds in some clerical workers.

Methods: Questionnaires were administered to 450 clerical workers between May and August 2000. The questionnaire included items concerning socio-demographic profile and noise-exposure history as well as use of mobile phone. The number of workers used in the final analysis was 394(Response rate 87.6%). Males numbered 308 and females 86. Hearing thresholds were measured in both ears using a pure-tone audiometer.

Results: In a univariate analysis, the duration and time of mobile phone use per day were significant in regards to hearing threshold at 1000Hz in males($p<0.05$). In a linear regression analysis with adjustment for age, education, military service, use of earphone, the group that used mobile phones for more than 49 months was more likely to have low hearing threshold at 1000Hz and 4000Hz in males($p<0.05$) as compared to the reference group(30). Use of a mobile phone for more than 60 minutes had a significant relation to hearing thresholds at 1000Hz in males($p<0.05$). However, in univariate and adjusted linear regression analyses, none of the mobile phone related variables were significant in females.

Conclusions: These results suggested that only certain elements of use of a mobile phone may be associated with hearing thresholds and there is a complex relationship between the use of mobile phone and hearing thresholds that may differ in terms of frequency (1000Hz and 4000Hz) and gender.

Key Words : Mobile Phone, Hearing Thresholds

70dB 68dB
(2001)

가 70dB

가 2001 1
2,674 56.6%가 가
(, 2001), 2004 3,100

가 1/3
(2000)

(, 2000).
1990 가

(Hermann & Hossmann, 1997; Hocking, 1998; Cox et al, 2000) 가 가

가 (Paredi et al, 2001). Kellenyi et al(1999) ABR(Auditory

800-1800MHz low power radio devices .
Brain Response) 20dB

DNA
(Lai & Singh, 1995; Szmigielski, 1996; Repacholi et al, 1997; Muscat et al, 2000)

가
가 ,

, EEG
(Mann & Roschke, 1996; Freude et al, 1998; Borbely et al, 1999; Preece et al, 1999; Hietanen et al, 2000; Koivisco et al, 2000; Krause et al, 2000).

1.

(Braune et al, 1998),
(Magras & Xenos, 1997),
(Royal Society of Canada, 1999; IEGMP, 2000)

가 450 가

(Naegeli et al., 1996; Hayes et al, 1997; Darmon et al, 1998; Kompis et al, 2000)

394 87.6%

2.

2000 6 1 2000 10 31 5
10

가 가 (Violanti and Marshall, 1996; Redelmeier, 1997; Dreyer et al, 1999)

가 (, ,)

,)
 (Beltone 112, USA)
 1000 4000Hz
 3. 21-60 49.8 % ,
 48.8 % 가 ,
 51.9 % , 37
 % , 41.9 % ,
 39.5 % 가
 56.7 % , 42.9
 t-test, % 가 (Table2).
 ANOVA ,
 1000 Hz
 4000 Hz 30 19.59±
 4.86, 21.7±13.41 가
 가 1000 Hz
 가 30 58.8 % , 1000 Hz 20 19.10±4.60
 가 20 83.7 % 가 , 가 61 17.3±5.55 가
 67.9 % , 72.1 % 가 , 4000 Hz
 가 39.3 % 가 가
 9.7 % . 1000 Hz 4000 Hz
 74.4 % , 44.2 % (Table 1). 가
 가
 31-48 50.2 % , 1000 Hz 4000 Hz
 30 52.3 % 가 . 19.85±3.9, 21.47±14.82 가

Table 1. General characteristics of the study population by sex

	Males (%) (N=308)	Females (%) (N=86)
Age		
20-29	92 (29.9)	72 (83.7)
30-39	181 (58.8)	14 (16.3)
40-	35 (11.4)	0 (0.0)
Educational attainment		
Less than high school	79 (25.6)	23 (26.7)
Graduated junior college	209 (67.9)	62 (72.1)
Graduated university	20 (6.5)	1 (1.2)
Military services		
Served, infantryman	212 (39.3)	-
Served, others	118 (38.3)	-
Civil defense	39 (12.7)	-
Exempted	30 (9.7)	-
Use of an earphone		
Yes	229 (74.4)	38 (44.2)
No	79 (25.6)	48 (55.8)

Table 2. Variables related to mobile phone use by sex

	Males (%) (N=308)	Females (%) (N=86)
Duration (month)		
- 30	73 (23.8)	45 (52.3)
31 - 48	154 (50.2)	33 (38.4)
49 -		
80 (26.1)	8 (9.3)	
Time per day (minute)		
- 20	100 (33.4)	18 (22.0)
21 - 60	149 (49.8)	40 (48.8)
61 -	50 (16.7)	24 (29.3)
Level of sound		
1 - 2 (small)	34 (11.0)	16 (18.6)
3	114 (37.0)	36 (41.9)
4 - 5 (loud)	160 (51.9)	34 (39.5)
Places used mostly		
Office	170 (56.7)	25 (29.8)
Car	63 (21.0)	11 (13.1)
House	17 (5.7)	36 (42.9)
Street & others	58 (18.8)	14 (16.3)

Table 3. Mean & standard deviation of mean hearing thresholds by use of mobile phone in males

	Mean 1K	P-value	Mean 4K	P-value
Duration (month)				
- 30	19.59 ± 4.86	0.019	21.47 ± 13.41	0.055
31 - 48	19.42 ± 4.42		21.56 ± 13.97	
49 -	17.81 ± 4.45		17.34 ± 11.78	
Time per day (minute)				
- 20	19.10 ± 4.60	0.012	20.75 ± 13.45	0.574
21 - 60	19.52 ± 4.12		20.87 ± 13.79	
61 -	17.30 ± 5.55		18.65 ± 11.50	
Level of sound				
1 - 2 (small)	18.24 ± 5.42	0.515	18.53 ± 13.42	0.119
3	19.07 ± 4.53		18.90 ± 11.32	
4 - 5 (large)	19.23 ± 4.44		21.95 ± 14.54	
Places used mostly				
Office	19.07 ± 4.78	0.894	21.24 ± 13.46	0.627
Car	18.89 ± 4.53		19.21 ± 12.24	
House	19.85 ± 3.90		21.47 ± 14.82	
Street & others	18.98 ± 4.32		19.18 ± 13.963)	

Table 4. Mean & standard deviation of mean hearing thresholds by use of mobile phone in females

	Mean 1K	P-value	Mean 4K	P-value
Duration (month)				
- 30	18.56 ± 4.78	0.312	13.50 ± 5.21	0.161
31 - 48	16.82 ± 5.09		11.29 ± 5.66	
49 -	18.13 ± 5.47		14.38 ± 7.41	
Time per day (minute)				
- 20	17.50 ± 4.77	0.940	12.78 ± 6.91	0.991
21 - 60	18.00 ± 4.81		12.63 ± 5.46	
61 -	17.92 ± 5.65		12.81 ± 5.58	
Level of sound				
1 - 2 (small)	19.22 ± 2.85	0.458	14.06 ± 4.46	0.479
3	17.36 ± 5.38		12.85 ± 6.19	
4 - 5 (large)	17.72 ± 5.31		11.99 ± 5.60	
Places used mostly				
Office	17.90 ± 4.82	0.861	12.90 ± 4.55	0.798
Car	16.82 ± 7.42		11.82 ± 5.93	
House	17.85 ± 4.52		12.43 ± 6.28	
Street & others	18.57 ± 4.46		13.93 ± 5.94	

(Table 3).

	Mean 1K	P-value	Mean 4K	P-value
1000Hz 4000Hz 31-48	16.82±	0.312	13.50 ± 5.21	0.161
5.09, 11.29±5.66	가 , 4000Hz		11.29 ± 5.66	
49	14.38±7.41		14.38 ± 7.41	
1000Hz 4000Hz				
4000Hz	1000Hz	0.940	1000Hz 4000Hz	0.991
, 4.043	가		0.324, 0.891,	
4000Hz	19.22		가	
±2.85, 14.06±4.46	가	4000Hz	1000Hz 4000Hz	가
	가	가	1.645, 2.484	

(Table5).

	Mean 1K	P-value	Mean 4K	P-value
5.94 가	18.57±4.46, 13.93±	0.861	12.90 ± 4.55	0.798
(Table4).	가		11.82 ± 5.93	
	가		12.43 ± 6.28	
1000Hz 4000Hz	1000Hz	0.940	1000Hz 4000Hz	0.991
30	31-48		4000Hz	
가	-0.087, -0.412		20	

Table 5. Multiple linear regression of mean hearing threshold (1K and 4K) after adjusting for age, educational attainments, military services and use of earphone in males

	1K			4K		
	*	SE() [†]	P-value	SE()	P-value	
Duration (month)						
31 - 48 vs - 30	-0.087	0.641	0.892	-0.412	1.878	0.826
49 - vs - 30	-1.821	0.740	0.014	-5.318	2.168	0.015
Time per day (minute)						
21 - 60 vs - 20	0.386	0.586	0.510	0.354	1.720	0.837
61 - vs - 20	-1.682	0.790	0.034	-1.679	2.318	0.469
Level of sound						
3 vs 1 - 2	0.743	0.885	0.402	0.891	2.565	0.729
4 - 5 vs 1 - 2	0.857	0.857	0.318	4.043	2.485	0.105
Places used mostly						
Car vs Office	-0.003	0.672	0.997	-2.692	1.958	0.170
House vs Office	1.645	1.189	0.168	2.484	3.464	0.474
Street & others vs Office	-0.235	0.694	0.735	-2.069	2.022	0.307

*regression coefficients, [†]standard errors of coefficients

Table 6. Multiple linear regression of mean hearing threshold (1K and 4K) after adjusting for age, educational attainments and use of earphone in females

	1K			4K		
	*	SE() [†]	P-value	SE()	P-value	
Duration (month)						
31 - 48 vs - 30	-1.704	1.171	0.150	-2.166	1.319	0.105
49 - vs - 30	-0.179	1.983	0.928	1.097	2.234	0.625
Time per day (minute)						
21 - 60 vs - 20	0.544	1.467	0.712	0.029	1.684	0.986
61 - vs - 20	0.481	1.636	0.769	0.292	1.879	0.877
Level of sound						
3 vs 1 - 2	-1.840	1.533	0.234	-1.225	1.742	0.484
4 - 5 vs 1 - 2	-1.493	1.537	0.334	-2.032	1.746	0.248
Places used mostly						
Car vs Office	-0.844	1.897	0.658	-0.787	2.151	0.715
House vs Office	0.129	1.410	0.927	-0.076	1.598	0.962
Street & others vs Office	0.840	1.745	0.632	1.359	1.979	0.494

1000Hz 4000Hz 21-60 가 0.840, 1.359 ,
 0.544, 0.029, 61 1000Hz 4000Hz,
 0.481, 0.292 4000Hz
 . 가 1000Hz
 4000Hz -1.840, -1.225, (Table6).
 -1.493, -2.032 .
 1000Hz 4000Hz

. Kellenyi et al(1999) 가 8
 ABR V 가
 (pulsed high-frequency EMF) V 가
 (Eulitz et al, 1998) 가 2-10 kHz
 가 20dB . 가
 가 , 가 가 가
 (radar-like pulsed RF 가 가
 field) 가 가
 (Kellenyi et al; 1999) .
 (Chou et al, 1985; UNEP/WHO/IRPA, 1993) 가
 4000Hz 1000Hz 가
 가 가
 1000Hz 1000Hz 4000Hz
 가 가 Kellenyi et al(1999)
 가 1000Hz
 , 4000Hz 4000Hz . Kellenyi et al(1999)
 가 p 0.055 가
 , , 가 ,
 4000Hz 1000Hz 가 ,
 49 30 가
 20 1000Hz 가
 가 60 audible acoustic vibra-
 tion
 가 .
 microwave hearing
 가 (IEGMP, 2000)
 , 가 가
 가 가
 4000Hz 가 1000Hz, 가

: 394

가

가

1000 400Hz

가

. Cox et al(2000)

)

:

100Hz

5-8 % 가

가

가

100Hz

400Hz

30

49

100Hz

20

60

(selec

가

tion bias) 가

가

가

:

가

가

100Hz

400Hz

가

가

. 2001

. 2001

가 . 2000

. 2000

Borbely AA. Huber R. Graf T. Fuchs B. Gallmann E. Achermann P. Pulsed high-frequency electromagnetic field affects human sleep and sleep electroencephalogram. *Neuroscience Letters* 1999; 275(3):207-10.

가

Braune S. Wrocklage C. Raczek J. Gailus T. Lucking CH. Resting blood pressure increase during exposure to a radio-frequency electromagnetic field. *Lancet* 1998; 351(9119):1857-8.

가

가

Chou CK. Yee KC. Guy AW. Auditory response in rats exposed to 2,450 MHz electromagnetic fields in a circularly polarized waveguide. *Bioelectromagnetics* 1985;6(3):323-6.

Cox, RAF, Luxon LM. Cerebral symptoms from mobile telephones. [Letter] *Occup Environ Med.* 2000;57(6):431.

Darmon P, Guillaume V, Wiart J, Dutour A, Oliver C.

- Do mobile cellular phones interfere with portable insulin pumps?. *Diabetes Care* 1998; 21: 1775.
- Dreyer NA, Loughlin JE, Rothman KJ. Cause-specific mortality in cellular telephone users. *JAMA* 1999;282(19): 1814-1816.
- IEGMP(Independent Exposure Group on Mobile Phones). Mobile phones and health. Report of an Independent Expert Group on Mobile Phones. UK Minister of public health. National Radiological Protection Board, Chilton, Oxon. 2000.
- Eulitz C, Ullsperger P, Freude G, Elbert T. Mobile phones modulate response patterns of human brain activity. *Neuroreport* 1998;9(1):3229-3232.
- Freude G, Ullsperger P, Eggert S, Ruppe I. Effects of microwaves emitted by cellular phones on human slow brain potentials. *Bioelectromagnetics* 1998;19:384-387.
- Hayes DL, Wang PJ, Reynolds DW, Estes M 3rd, Griffith JL, Steffens RA, Carlo GL, Findlay GK, Johnson CM. Interference with cardiac pacemakers by cellular telephones. *NEJM* 1997; 336(21):1473-9.
- Hermann DM, Hossmann KA. Neurological effects of microwave exposure related to mobile communication. *J Neurol Sci* 1997;152:1-14.
- Hietanen M, Kovala T, Hamalainen AM. Human brain activity during exposure to radiofrequency fields emitted by cellular phones [see comments]. *Scand J Work, Environ and Health* 2000; 26(2):87-92.
- Hocking B. Preliminary report: symptoms associated with mobile phone use. *Occup Med.* 1998; 48(6):357-60.
- Hyland, GJ. Physics and biology of mobile telephony. *Lancet* 2000;356(9244): 1833-1836.
- Kellenyi L, Thuroczy GY, Faludy B, Lenard L. Effects of mobile GSM radiotelephone exposure on the auditory brainstem response. *Neurobiology* 1999; 7(1): 79-81.
- Kosvisto M, Revonsuo A, Krause CM, Harala C, Sillanmaki L, Laine M, Hamalainen H. Effects of 900MHz EMF emitted by cellular phones on response times in humans. *NeuroReport* 2000; 11: 1641-1643.
- Kompis M, Negri S, Hausler R. Electromagnetic interference of bone-anchored hearing aids by cellular phones. *Acta Otolaryngol* 2000; 120: 855-859.
- Krause CM, Sillanmaki L, Koivisto M, Haggqvist A, Saarela C, Revonsuo A, Laine M, Hamalainen H. Effects of EMF emitted by cellular phones on the EEG during a memory task. *Neuroreport* 2000; 11(4): 761-764.
- Lai H, Singh NP. Acute low-intensity microwave exposure increases DNA single-strand breaks in rat brain cells. *Bioelectromagnetics* 1995;16(3):207-10.
- Magras IN, Xenos TD. RF radiation-induced changes in the prenatal development of mice. *Bioelectromagnetics* 1997;18(6):455-461.
- Mann K, Roschke J. Effects of pulsed high-frequency electromagnetic fields on human sleep. *Neuropsychobiology* 1996; 33: 41-47.
- Muscat JE, Malkin MG, Thompson S, Shore RE, Stellman SD, McRee D, Neugut AI, Wynder EL. Handheld cellular telephone use and risk of brain cancer. *JAMA* 284(23):3001-7.
- Naegeli B, Osswald S, Deola M, Burkart F. Intermittent pacemaker dysfunction caused by digital mobile telephones. *Electrophysiol* 1996; 27(6):1471-1477.
- Paredi P, Kharitonov SA, Hanazawa T, Barnes PJ. Local vasodilator response to mobile phones. *Laryngoscope* 2001; 111(1): 159-62.
- Preece AW, Iwi G, Davies-Smith A, Wesnes K, Butler S, Lim E, Valey A. Effect of a 915-MHz simulated mobile phone signal on cognitive function in man. *Int J Radiat Biol.* 1999;75(4):447-56.
- Royal Society of Canada. A review of the potential health risks of radiofrequency fields from wireless telecommunication devices. 1999 Ottawa, Ontario.
- Redelmeier DA, Tibshirani RJ. Association between cellular-telephone calls and motor vehicle collisions. *NEJM* 1997; 336(7): 453-458.
- Repacholi MH, Basten A, Gebiski V, Noonan D, Finnie J, Harris AW. Lymphomas in EμPim1 transgenic mice exposed to pulsed 900 mHz electromagnetic fields. *Radiation Res* 1997; 147: 631-640.
- Szmigielski S. Cancer morbidity in subjects occupationally exposed to high frequency(RF & microwave) EM radiation. *Sci Total Environ* 1996; 180(11): 9-17.
- UNEP/WHO/IRPA. Electromagnetic fields(300Hz-300GHz). 1993 Geneva, World Health Organization, Environmental Health Criteria 137
- Van Leeuwen GMJ, Lagendijk JJW, Van

- Leersum BIAM, Zwamborn APM, Hornssleuth SN, Kotte ANT. Calculation of brain temperature due to exposure to a mobile phone. *Phys. Med. Biol.* 44 2367 1999.
- Violanti JM, Marshall JR. Cellular phones and traffic accidents: an epidemiological approach. *Accid Anal Prev* 1996;28(2):265-170.
- WHO, 2000. Fact sheet on electromagnetic fields and public health: mobile phones and their base stations. World Health Organization, Geneva.